

t7_scmring1

(TMTL6eqcQ8ERNdXa39B4VSbepmRxjMqFhaN)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_card_3 : \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_ami_2 : \iota$ be given. Let $k1_scmring1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_scmring1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_afinsq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_ami_2 : \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. k1_funct_1 (k16_funcop_1 X0 X1) X0 = X1 \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 \\ & (k1_funct_4 X2 X1) X0 = k1_funct_1 X1 X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k6_afinsq_1 X0 X1)) \wedge (v1_funct_1 (k6_afinsq_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (v4_funct_1 (k4_card_3 X0)) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k3_relat_1 X0 X1)) \wedge (v1_funct_1 (k3_relat_1 X0 X1))) \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k2_funcop_1 X0 X1)) \wedge (v1_funct_1 (k2_funcop_1 X0 X1)) \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k3_relat_1 X0 X1) \quad (9)$$

Assume the following.

$$(v1_funct_1 k3_ami_2) \wedge ((v1_funct_2 k3_ami_2 k1_ami_2 np_2) \wedge (m1_subset_1 k3_ami_2 (k1_zfmisc_1 (k2_zfmisc_1 k1_ami_2 np_2)))) \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarski X0) X1 \quad (11)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (k4_card_3 (k3_relat_1 k3_ami_2 (k1_scmring1 X0)))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (k3_scmring1 X0 X1 X2 = k1_funct_4 X1 (k16_funcop_1 k5_numbers X2)))) \quad (12)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (k1_scmring1 X0 = k6_afinsq_1 k5_numbers (u1_struct_0 X0)) \quad (13)$$

Assume the following.

$$\forall X0. \forall X1. k2_funcop_1 X0 X1 = k2_zfmisc_1 X0 (k1_tarski X1) \quad (14)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (15)$$

Assume the following.

$$\forall X0.(v4_funct_1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v1_relat_1 X1) \wedge (v1_funct_1 X1)) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (17)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\forall X1.(m1_subset_1 X1 (k4_card_3 (k3_relat_1 k3_ami_2 (k1_scmring1 X0)))) \Rightarrow (\forall X2.(v7_ordinal1 X2) \Rightarrow (k1_funct_1 (k3_scmring1 X0 X1 X2) k5_numbers = X2)))$$