

t7_seqm_3

(TMaUSGTRFzN4Jibs1Xy5MChno9nEz7QJ3sr)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v8_valued_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\
 & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\
 & (((\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (r1_xxreal_0 \\
 & (k8_nat_1 k1_numbers X0 (k2_nat_1 X1 np_1)) (k8_nat_1 k1_numbers \\
 & X0 X1))) \Rightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\
 & (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (r1_xxreal_0 \\
 & (k8_nat_1 k1_numbers X0 (k2_nat_1 X1 X2)) (k8_nat_1 k1_numbers \\
 & X0 X1)))))) \wedge (((\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\
 & (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (r1_xxreal_0 \\
 & (k8_nat_1 k1_numbers X0 (k2_nat_1 X1 X2)) (k8_nat_1 k1_numbers \\
 & X0 X1)))))) \Rightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\
 & (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow ((r1_xxreal_0 \\
 & X1 X2) \Rightarrow (r1_xxreal_0 (k8_nat_1 k1_numbers X0 X2) (k8_nat_1 k1_numbers \\
 & X0 X1)))))) \wedge (((\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\
 & (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow ((r1_xxreal_0 \\
 & X1 X2) \Rightarrow (r1_xxreal_0 (k8_nat_1 k1_numbers X0 X2) (k8_nat_1 k1_numbers \\
 & X0 X1)))))) \Rightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\
 & (r1_xxreal_0 (k8_nat_1 k1_numbers X0 (k2_nat_1 X1 np_1)) (k8_nat_1 \\
 & k1_numbers X0 X1))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\
& (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\
& ((v8_valued_0 X0) \Leftrightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\
& (r1_xxreal_0 (k8_nat_1 k1_numbers X0 (k2_nat_1 X1 np_1)) (k8_nat_1 \\
& k1_numbers X0 X1))))
\end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\
& (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\
& ((v8_valued_0 X0) \Leftrightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\
& (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (r1_xxreal_0 \\
& (k8_nat_1 k1_numbers X0 (k2_nat_1 X1 X2)) (k8_nat_1 k1_numbers \\
& X0 X1))))))
\end{aligned}$$