

## t7\_sin\_cos7

(TMa5LDnZJ8FYjZcbwp2GGAX4EAb98N2rJ3R)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k12\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_0 : \iota$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $k7\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 X1) \wedge (\neg v3\_xxreal\_0 X1) \wedge (\neg v2\_xxreal\_0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2.(v1\_xreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Leftrightarrow (r1\_xxreal\_0 (k2\_xcmplx\_0 X0 X2) (k2\_xcmplx\_0 X1 X2)))))) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (v3\_xxreal\_0 X1)) \Rightarrow (v3\_xxreal\_0 X0))) \quad (4)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (5)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (6)$$

Assume the following.

$$k2\_xcmplx\_0 \text{ np\_}1 (k4\_xcmplx\_0 \text{ np\_}1) = \text{np\_}0 \quad (7)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k12\_binop\_2 X0 X1 = k7\_xcmplx\_0 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k10\_binop\_2 X0 X1 = k6\_xcmplx\_0 X0 X1) \quad (10)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0)\wedge((v1\_xcmplx\_0 X0)\wedge((v1\_xxreal\_0 X0)\wedge(v1\_xreal\_0 X0))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(v1\_xreal\_0 (k7\_xcmplx\_0 X0 X1)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(v1\_xreal\_0 (k2\_xcmplx\_0 X0 X1)) \quad (13)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow((v1\_xcmplx\_0 (k4\_xcmplx\_0 X0))\wedge(v1\_xreal\_0 (k4\_xcmplx\_0 X0))) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.(((\neg v3\_xxreal\_0 X0)\wedge(v1\_xreal\_0 X0))\wedge((\neg v3\_xxreal\_0 X1)\wedge(v1\_xreal\_0 X1)))\Rightarrow(\neg v3\_xxreal\_0 (k7\_xcmplx\_0 X0 X1)) \quad (15)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0)\Rightarrow(\forall X1.(v1\_xcmplx\_0 X1)\Rightarrow(k6\_xcmplx\_0 X0 X1 = k2\_xcmplx\_0 X0 (k4\_xcmplx\_0 X1))) \quad (16)$$

Assume the following.

$$\forall X0.((v1\_xxreal\_0 X0)\wedge(v3\_xxreal\_0 X0))\Rightarrow((\neg v1\_xboole\_0 X0)\wedge((v1\_xreal\_0 X0)\wedge(\neg v2\_xxreal\_0 X0))) \quad (17)$$

Assume the following.

$$\forall X0.((v1\_xxreal\_0 X0) \wedge (v2\_xxreal\_0 X0)) \Rightarrow ((\neg v1\_xboole\_0 X0) \wedge ((v1\_xxreal\_0 X0) \wedge (\neg v3\_xxreal\_0 X0))) \quad (18)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (19)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (20)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 np\_1 X1)) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k12\_binop\_2 (k10\_binop\_2 X1 np\_1) X0))))$$