

t7_wsierp_1 (TMKoEVWjDLN- MeGw1guCNvMsqbG2pQWoMZi1)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k3_int_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_int_1 X0) \Rightarrow (\forall X1.(v1_int_1 X1) \Rightarrow (\forall X2. \\ & (v1_int_1 X2) \Rightarrow ((k3_int_2 X0 X1 = np_1) \wedge (k3_int_2 X2 X1 = np_1)) \Rightarrow \\ & (k3_int_2 (k3_xcmplx_0 X0 X2) X1 = np_1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_int_1 X0) \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7_ordinal1 X2) \Rightarrow ((k3_int_2 X0 X1 = np_1) \wedge (k3_int_2 X2 X1 = np_1)) \Rightarrow \\ & (k3_int_2 (k3_xcmplx_0 X0 X2) X1 = np_1)))) \end{aligned}$$