

t7\_zfmisc\_1 (TMTFk-  
fJp6DYPNrpoh5rgBDv72CTBDS7hGC4)

October 27, 2020

Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k2\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2\_tarski X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 = X0) \vee (X3 = X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1\_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (3)$$

**Theorem 1**  $\forall X0. \forall X1. r1\_tarski (k1\_tarski X0) (k2\_tarski X0 X1).$