

t81_funcop_1 (TM-
FZVvG8S2YSRSAj83ePeMwxpexALPdpbWr)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))) \Rightarrow (r1_tarski (k7_relset_1 X0 X1 X3 X2) X1) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (v1_funct_1 (k7_funcop_1 X0 X1)) \wedge ((v1_funct_2 \\ & (k7_funcop_1 X0 X1) X0 (k1_tarski X1)) \wedge (m1_subset_1 (k7_funcop_1 \\ & X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarski X1)))) \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. r1_tarski (k7_relset_1 X0 (k1_tarski \\ & X2) (k7_funcop_1 X0 X2) X1) (k1_tarski X2) \end{aligned}$$