

t81_qc_lang2
 $(TMXPXrgFz4TSuHshua5pnWhxXdhuTCsop)$

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $r2_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_qc_lang2 : \iota \Rightarrow \iota$ be given. Let $k12_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $r3_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) &\Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((r2_qc_lang2 X0 X1 (k12_qc_lang1 X0)) \Leftrightarrow (X1 = k12_qc_lang1 \\ X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) &\Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((r3_qc_lang2 X0 X1 (k1_qc_lang2 X0)) \Leftrightarrow (r2_qc_lang2 X0 X1 \\ (k12_qc_lang1 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0) \wedge ((m1_subset_1 \\ X1 (k9_qc_lang1 X0)) \wedge (m1_subset_1 X2 (k9_qc_lang1 X0)))) &\Rightarrow (r2_qc_lang2 \\ X0 X1 X1) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k1_qc_lang2 X0) (k9_qc_lang1 \\ X0)) \quad (4)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k12_qc_lang1 X0) (\\ k9_qc_lang1 X0)) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) &\Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0)) \Rightarrow ((r3_qc_lang2 \\ X0 X1 X2) \Leftrightarrow ((r2_qc_lang2 X0 X1 X2) \wedge (X1 \neq X2))))) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1\ X0)) \Rightarrow ((r2_qc_lang2\ X0\ X1\ (k1_qc_lang2\ X0)) \Leftrightarrow ((X1 = k1_qc_lang2\ X0) \vee (X1 = k12_qc_lang1\ X0)))) \end{aligned}$$