

t81_xxreal_3
(TMP31C8fYf9hgvJ6GuX767CVLHJCNsFKger)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k4_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Let $k2_xxreal_0 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 np_1 X0 = X0) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\neg(\neg X0 \in k1_numbers) \wedge ((X0 \neq k1_xxreal_0) \wedge (X0 \neq k2_xxreal_0))) \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (4)$$

Assume the following.

$$v3_xxreal_0 k2_xxreal_0 \quad (5)$$

Assume the following.

$$v2_xxreal_0 k1_xxreal_0 \quad (6)$$

Assume the following.

$$\begin{aligned}
& \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\
& (v1_xxreal_0 X2) \Rightarrow (((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow ((X2 = \\
& k4_xxreal_3 X0 X1) \Leftrightarrow (\exists X3.(v1_xcmplx_0 X3) \wedge (\exists X4. \\
& (v1_xcmplx_0 X4) \wedge ((X0 = X3) \wedge ((X1 = X4) \wedge (X2 = k3_xcmplx_0 X3 X4)))))) \wedge \\
& ((\neg(\neg(v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \wedge (((v2_xxreal_0 X0) \wedge \\
& (v2_xxreal_0 X1)) \vee ((v3_xxreal_0 X0) \wedge (v3_xxreal_0 X1))) \wedge (\neg \\
& X2 = k4_xxreal_3 X0 X1) \Leftrightarrow (X2 = k1_xxreal_0))) \wedge (\neg(\neg(v1_xreal_0 \\
& X0) \wedge (v1_xreal_0 X1)) \wedge (((v2_xxreal_0 X0) \wedge (v3_xxreal_0 X1)) \vee \\
& ((v3_xxreal_0 X0) \wedge (v2_xxreal_0 X1))) \wedge (\neg(X2 = k4_xxreal_3 X0 X1) \Leftrightarrow \\
& (X2 = k2_xxreal_0))) \wedge (\neg(\neg(v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \wedge \\
& ((\neg(\neg(v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \wedge (((v2_xxreal_0 X0) \wedge (\\
& v2_xxreal_0 X1)) \vee ((v3_xxreal_0 X0) \wedge (v3_xxreal_0 X1)))) \wedge (\neg \\
& (\neg(v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \wedge (((v2_xxreal_0 X0) \wedge (v3_xxreal_0 \\
& X1)) \vee ((v3_xxreal_0 X0) \wedge (v2_xxreal_0 X1)))) \wedge (\neg(X2 = k4_xxreal_3 \\
& X0 X1) \Leftrightarrow (X2 = k6_numbers)))))))))
\end{aligned} \tag{7}$$

Assume the following.

$$k1_xxreal_0 = k1_numbers \tag{8}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \tag{9}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \tag{10}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \tag{11}$$

Theorem 1 $\forall X0.(v1_xxreal_0 X0) \Rightarrow (k4_xxreal_3 \text{ np}_{-1} X0 = X0).$