

t82_funct_1

(TMXZWq1DH5FbHGozxroCM6FQhVnoQfHQVZX)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X2 = k8_relat_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in k9_xtuple_0 \\ & X0) \wedge (k1_funct_1 X0 X3 \in X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X2 = k7_relat_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow (\exists X4. (X4 \in k9_xtuple_0 \\ & X0) \wedge ((X4 \in X1) \wedge (X3 = k1_funct_1 X0 X4))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v2_funct_1 X0) \Leftrightarrow \\ & (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \wedge ((X2 \in k9_xtuple_0 \\ & X0) \wedge (k1_funct_1 X0 X1 = k1_funct_1 X0 X2))) \Rightarrow (X1 = X2))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow \\ & (X2 \in X1)) \end{aligned} \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((v2_funct_1 \\ & X1) \Rightarrow (r1_tarski (k8_relat_1 X1 (k7_relat_1 X1 X0)) X0)) \end{aligned}$$