

t82_xxreal_1 (TMF-
PbnHx9kYvTVZ1ABKJTXyksX8ZLpJLmKf)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(r1_tarski X0 X1) \wedge (r2_xboole_0 X1 X0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3. (v1_xxreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_tarski (k1_xxreal_1 X1 X2) (k1_xxreal_1 \\ & X0 X3)))))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3. (v1_xxreal_0 X3) \Rightarrow (\neg(r1_xxreal_0 \\ & X0 X1) \wedge ((r2_xboole_0 (k1_xxreal_1 X0 X1) (k1_xxreal_1 X2 X3)) \wedge \\ & ((r1_xxreal_0 X0 X2) \wedge (r1_xxreal_0 X3 X1)))))) \end{aligned}$$