

t83_setlim_2 (TMbGK- CLZk4RHdLVkU2Z138y1JUAXUERtgeN)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_kurato_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_setlim_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (\forall X2. (X2 \in k4_kurato_0 \\ & X0 X1) \Leftrightarrow (\forall X3. (m1_subset_1 X3 k5_numbers) \Rightarrow (\exists X4. (\\ & m1_subset_1 X4 k5_numbers) \wedge (X2 \in k3_funct_2 k5_numbers (k9_setfam_1 \\ & X0) X1 (k2_nat_1 X3 X4)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.k9_setfam.1 X0 = k1_zfmisc.1 X0 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((v1_funct.1 X1)\wedge((v1_funct.2 X1 k5_numbers X0)\wedge(m1_subset.1 X1 (k1_zfmisc.1 (k2_zfmisc.1 k5_numbers X0))))))\wedge(v7_ordinal1 X2))\Rightarrow(k8_nat.1 X0 X1 X2 = k1_funct.1 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset.1 X1 (k1_zfmisc.1 X0))\Rightarrow(k7_subset.1 X0 X1 X2 = k4_xboole.0 X1 X2) \quad (7)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole.0 X0)\wedge(((v1_funct.1 X2)\wedge((v1_funct.2 X2 X0 X1)\wedge(m1_subset.1 X2 (k1_zfmisc.1 (k2_zfmisc.1 X0 X1))))))\wedge(m1_subset.1 X3 X0)))\Rightarrow(k3_funct.2 X0 X1 X2 X3 = k1_funct.1 X2 X3) \quad (9)$$

Assume the following.

$$\forall X0.(\neg v1_xboole.0 X0)\Rightarrow(\exists X1.(m1_subset.1 X1 (k1_zfmisc.1 X0))\wedge(\neg v1_xboole.0 X1)) \quad (10)$$

Assume the following.

$$(\neg v1_xboole.0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole.0 X0)\wedge((\neg v1_xboole.0 X1)\wedge(m1_subset.1 X1 (k1_zfmisc.1 X0))))\Rightarrow(\exists X2.m2_subset.1 X2 X0 X1) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole.0 X0)\wedge((\neg v1_xboole.0 X1)\wedge(m1_subset.1 X1 (k1_zfmisc.1 X0))))\Rightarrow(\forall X2.(m2_subset.1 X2 X0 X1)\Rightarrow(m1_subset.1 X2 X0)) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((v1_funct_1 X1)\wedge((v1_funct_2 \\ & X1 k5_numbers (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0))))))\wedge(m1_subset_1 \\ & X2 (k1_zfmisc_1 X0)))\Rightarrow((v1_funct_1 (k7_setlim_2 X0 X1 X2))\wedge((\\ & v1_funct_2 (k7_setlim_2 X0 X1 X2) k5_numbers (k9_setfam_1 X0))\wedge \\ & (m1_subset_1 (k7_setlim_2 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \end{aligned} \quad (14)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 \\ & X1))\Rightarrow(m2_subset_1 (k2_nat_1 X0 X1) k1_numbers k5_numbers) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0))))))\Rightarrow(\forall X2.(m1_subset_1 X2 \\ & (k1_zfmisc_1 X0))\Rightarrow(\forall X3.((v1_funct_1 X3)\wedge((v1_funct_2 \\ & X3 k5_numbers (k9_setfam_1 X0))\wedge(m1_subset_1 X3 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0))))))\Rightarrow((X3 = k7_setlim_2 \\ & X0 X1 X2)\Leftrightarrow(\forall X4.(m1_subset_1 X4 k5_numbers)\Rightarrow(k8_nat_1 (\\ & k9_setfam_1 X0) X3 X4 = k7_subset_1 X0 X2 (k8_nat_1 (k9_setfam_1 \\ & X0) X1 X4)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(X2 = k4_xboole_0 X0 X1)\Leftrightarrow(\forall X3. \\ & (X3 \in X2)\Leftrightarrow((X3 \in X0)\wedge(\neg X3 \in X1))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(r1_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow \\ & (X2 \in X1)) \end{aligned} \quad (19)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 \\ & X1))\Rightarrow(k2_nat_1 X0 X1 = k2_nat_1 X1 X0) \end{aligned} \quad (20)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (21)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 k5_numbers (k9_setfam_1 X0)) \wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 \\ & X0)))))) \Rightarrow (r1_tarski (k7_subset_1 X0 X1 (k4_kurato_0 X0 X2)) (k4_kurato_0 \\ & X0 (k7_setlim_2 X0 X2 X1)))) \end{aligned}$$