

t84_funcop_1
 (TMJ7GP92owaNjKJdGznzSr7e4snappjJ8NT)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((r1_tarski (k16_funcop_1 X0 X1) X2) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (k1_funct_1 X2 X0 = X1))) \tag{1}$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X0 \in k9_xtuple_0 X1) \Rightarrow (r1_tarski (k16_funcop_1 X0 (k1_funct_1 X1 X0)) X1))$$