

t84_zfmisc_1 (TMGnE-
FCxLttZdYukhE8hMg5AycW7HhRfYqX)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_zfmisc_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow (\exists X4. \exists X5. (X4 \in X0) \wedge ((X5 \in X1) \wedge (X3 = k4_tarski X4 X5)))) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. \forall X3. \neg (r1_tarski X0 (k2_zfmisc_1 X1 X2)) \wedge ((X3 \in X0) \wedge (\forall X4. \forall X5. \neg (X4 \in X1) \wedge ((X5 \in X2) \wedge (X3 = k4_tarski X4 X5))))$$