

t85_quaterni (TMdyNd- ddpqY9wP4DCDoHt3k9YJweCvymD)

October 27, 2020

Let $v1_quaterni : \iota \Rightarrow o$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k32_quaterni : \iota \Rightarrow \iota$ be given. Let $k29_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k26_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_quaterni : \iota$ be given. Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (\forall X1.(v1_quaterni X1) \Rightarrow (r1_xreal_0 (k32_quaterni (k26_quaterni X0 X1)) (k2_xcmplx_0 (k32_quaterni X0) (k32_quaterni X1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (\forall X1.(v1_quaterni X1) \Rightarrow (\forall X2.(v1_quaterni X2) \Rightarrow (k29_quaterni X1 X2 = k26_quaterni (k29_quaterni X1 X0) (k29_quaterni X0 X2)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_quaterni X0) \wedge (v1_quaterni X1)) \Rightarrow (m1_subset_1 (k29_quaterni X0 X1) k1_quaterni) \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_quaterni) \Rightarrow (v1_quaterni X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_quaterni X0) \Rightarrow (\forall X1.(v1_quaterni X1) \Rightarrow (\forall X2.(v1_quaterni X2) \Rightarrow (r1_xreal_0 (k32_quaterni (k29_quaterni X0 X1)) (k2_xcmplx_0 (k32_quaterni (k29_quaterni X0 X2)) (k32_quaterni (k29_quaterni X2 X1))))))$$