

t85_setlim_2

(TMZRv44QpAXu8vzUXV2CFxgMtisakPwhXfu)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_kurato_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_setlim_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\
 & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 k5_numbers (k9_setfam_1 X0)) \wedge (m1_subset_1 X2 \\
 & (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0)))))) \Rightarrow \\
 & (\forall X3. (m1_subset_1 X3 (k1_zfmisc_1 X0)) \Rightarrow ((\forall X4. (\\
 & m1_subset_1 X4 k5_numbers) \Rightarrow (k3_funct_2 k5_numbers (k9_setfam_1 \\
 & X0) X2 X4 = k5_subset_1 X0 X3 (k3_funct_2 k5_numbers (k9_setfam_1 \\
 & X0) X1 X4))) \Rightarrow (r1_tarski (k5_subset_1 X0 X3 (k4_kurato_0 X0 X1)) \\
 & (k4_kurato_0 X0 X2))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. k9_setfam_1 X0 = k1_zfmisc_1 X0 \tag{2}$$

Assume the following.

$$\begin{aligned}
 & \forall X0. \forall X1. \forall X2. (((v1_funct_1 X1) \wedge ((v1_funct_2 \\
 & X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\
 & X0)))))) \wedge (v7_ordinal1 X2)) \Rightarrow (k8_nat_1 X0 X1 X2 = k1_funct_1 X1 X2)
 \end{aligned} \tag{3}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ & (((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))))))\wedge(m1_subset_1 X3 X0)))\Rightarrow(k3_funct_2 X0 \\ & X1 X2 X3 = k1_funct_1 X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((v1_funct_1 X1)\wedge((v1_funct_2 \\ & X1 k5_numbers (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0))))))\wedge(m1_subset_1 \\ & X2 (k1_zfmisc_1 X0)))\Rightarrow((v1_funct_1 (k9_setlim_2 X0 X1 X2))\wedge((\\ & v1_funct_2 (k9_setlim_2 X0 X1 X2) k5_numbers (k9_setfam_1 X0))\wedge \\ & (m1_subset_1 (k9_setlim_2 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0))))))\Rightarrow(\forall X2.(m1_subset_1 X2 \\ & (k1_zfmisc_1 X0))\Rightarrow(\forall X3.((v1_funct_1 X3)\wedge((v1_funct_2 \\ & X3 k5_numbers (k9_setfam_1 X0))\wedge(m1_subset_1 X3 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0))))))\Rightarrow((X3 = k9_setlim_2 \\ & X0 X1 X2)\Leftrightarrow(\forall X4.(m1_subset_1 X4 k5_numbers)\Rightarrow(k8_nat_1 (\\ & k9_setfam_1 X0) X3 X4 = k5_subset_1 X0 X2 (k8_nat_1 (k9_setfam_1 \\ & X0) X1 X4)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(\forall X2. \\ & ((v1_funct_1 X2)\wedge((v1_funct_2 X2 k5_numbers (k9_setfam_1 X0))\wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 \\ & X0))))))\Rightarrow(r1_tarski (k5_subset_1 X0 X1 (k4_kurato_0 X0 X2)) (k4_kurato_0 \\ & X0 (k9_setlim_2 X0 X2 X1)))) \end{aligned}$$