

t85_xboole_1 (TMS-
dPmwdX6RRDb6oiuq4CjB8NMinDWnqxcm)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (\neg(\neg r1_xboole_0 X0 X1) \wedge (\forall X2. \neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2. (X2 \in X0) \wedge (X2 \in X1))) \wedge (r1_xboole_0 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (r1_tarski X0 X1) \Rightarrow (r1_xboole_0 X0 (k4_xboole_0 X2 X1))$$