

t85_xboolean
(TMb2GYsq5Mk9NP8B9a9Gq6eMAvVBDS8UfWt)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k6_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k5_xboolean X0 (k10_xboolean (k3_xboolean X0) X1) = k5_xboolean X0 (k3_xboolean X1))) \tag{1}$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (k3_xboolean (k3_xboolean X0) = X0) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k10_xboolean X0 X1)) \tag{3}$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (v1_xboolean (k3_xboolean X0)) \tag{4}$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k6_xboolean X0 X1 = k5_xboolean (k3_xboolean X0) X1)) \tag{5}$$

Theorem 1

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k6_xboolean X0 (k10_xboolean X0 X1) = k5_xboolean (k3_xboolean X0) (k3_xboolean X1)))$$