

t86_funct_2 (TM-
bLQPTcRTzcdWA6CUXnzMzDKJXb486CP1n)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow \\ & (((X1 = k1_xboole_0) \Rightarrow (X0 = k1_xboole_0)) \Rightarrow (k5_partfun1 X0 X1 (k3_partfun1 \\ & X2 X0 X1) = k1_tarski X2)) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 X0) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 X0)))))) \Rightarrow (k5_partfun1 \\ & X0 X0 (k3_partfun1 X1 X0 X0) = k1_tarski X1) \end{aligned}$$