

t86_newton (TMZGvdMQTypxUFSsxyGzy-
bGy2Rid1MTZW9K)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (r1_xxreal_0 (k1_nat_1 X0 np_1) (k2_newton np_2 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_newton X0 X1 = k1_newton X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (v7_ordinal1 (k1_newton X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Theorem 1 $\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg r1_xxreal_0 (k2_newton np_2 X0) X0)$.