

t87\_flang\_2  
(TMNyCuxxxubJHajUYf3FC2z1QdYu4daQa2b)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k4\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_catalan2 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k4\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \neg(v1\_xboole\_0 X0) \wedge ((X0 \neq X1) \wedge (v1\_xboole\_0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (k2\_flang\_2 X0 X1 = k4\_subset\_1 (k8\_afinsq\_1 X0) (k4\_flang\_1 X0 (k2\_flang\_1 X0)) X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. k2\_xboole\_0 X0 (k3\_xboole\_0 X1 X2) = k3\_xboole\_0 (k2\_xboole\_0 X0 X1) (k2\_xboole\_0 X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow (k9\_subset\_1 X0 X1 X2 = k3\_xboole\_0 X1 X2) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0))) \Rightarrow (k4\_subset\_1 X0 X1 X2 = k2\_xboole\_0 X1 X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X1 (k3\_catalan2 X0))\Rightarrow(k4\_flang\_1 X0 X1 = k1\_tarSKI X1) \quad (6)$$

Assume the following.

$$\forall X0.k3\_catalan2 X0 = k8\_afinsq\_1 X0 \quad (7)$$

Assume the following.

$$\forall X0.k2\_flang\_1 X0 = k4\_afinsq\_1 X0 \quad (8)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0)\wedge(v1\_xreal\_0 X0) \quad (9)$$

Assume the following.

$$\exists X0.v1\_xboole\_0 X0 \quad (10)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k4\_afinsq\_1 X0))\wedge((v5\_relat\_1 (k4\_afinsq\_1 X0) X0)\wedge((v5\_ordinal1 (k4\_afinsq\_1 X0))\wedge((v1\_funct\_1 (k4\_afinsq\_1 X0))\wedge((v1\_xboole\_0 (k4\_afinsq\_1 X0))\wedge(v1\_finset\_1 (k4\_afinsq\_1 X0)))))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 X0))\Rightarrow(m1\_subset\_1 (k9\_subset\_1 X0 X1 X2) (k1\_zfmisc\_1 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X1 (k3\_catalan2 X0))\Rightarrow(m1\_subset\_1 (k4\_flang\_1 X0 X1) (k1\_zfmisc\_1 (k3\_catalan2 X0))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0)))\Rightarrow(m1\_subset\_1 (k2\_flang\_2 X0 X1) (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \quad (14)$$

Assume the following.

$$\forall X0.m1\_subset\_1 (k2\_flang\_1 X0) (k3\_catalan2 X0) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 X0))\Rightarrow(k9\_subset\_1 X0 X1 X2 = k9\_subset\_1 X0 X2 X1) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.k3\_xboole\_0 X0 X1 = k3\_xboole\_0 X1 X0 \quad (17)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 \\ & X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k8\_afinsq\_1 \\ X0))) \Rightarrow (k2\_flang\_2 X0 (k9\_subset\_1 (k8\_afinsq\_1 X0) X1 X2) = k9\_subset\_1 \\ & (k8\_afinsq\_1 X0) (k2\_flang\_2 X0 X1) (k2\_flang\_2 X0 X2))) \end{aligned}$$