

t87_funct_1
(TMH3cdP9ZZAF5h1wdbYPw5tPCnxu8QVqfs1)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ (X2 = k7_relat_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow (\exists X4. (X4 \in k9_xtuple_0 \\ X0) \wedge ((X4 \in X1) \wedge (X3 = k1_funct_1 X0 X4)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v2_funct_1 X0) \Leftrightarrow \\ (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \wedge ((X2 \in k9_xtuple_0 \\ X0) \wedge (k1_funct_1 X0 X1 = k1_funct_1 X0 X2)))) \Rightarrow (X1 = X2))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{4}$$

Theorem 1

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 \\ X2)) \Rightarrow (((r1_tarski (k7_relat_1 X2 X0) (k7_relat_1 X2 X1)) \wedge ((r1_tarski \\ X0 (k9_xtuple_0 X2)) \wedge (v2_funct_1 X2))) \Rightarrow (r1_tarski X0 X1)) \end{aligned}$$