

t89_funct_7

(TMbUiwfAnJMi4tULqsyKeiyqkZGairY7TVt)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & \forall X3.(X2 \neq X3) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X3 = k1_funct_1 \\ & \quad X0 X3)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & k9_xtuple_0 (k2_funct_7 X0 X2 X1) = k9_xtuple_0 X0) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((\\ & v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski X0 X1) \Leftrightarrow ((r1_tarski \\ & (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \wedge (\forall X2.(X2 \in k9_xtuple_0 \\ & X0) \Rightarrow (k1_funct_1 X0 X2 = k1_funct_1 X1 X2)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1_relat_1 X0) \wedge (v1_funct_1 \\ & X0)) \Rightarrow ((v1_relat_1 (k2_funct_7 X0 X1 X2)) \wedge (v1_funct_1 (k2_funct_7 \\ & \quad X0 X1 X2))) \end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((\\ & v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \forall X3.(r1_tarski \\ & X1 X0) \Rightarrow ((X2 \in k9_xtuple_0 X1) \vee (r1_tarski X1 (k2_funct_7 X0 X2 X3)))))) \end{aligned}$$