

t89_xboolean
(TMXA1EokyaWDUwXu8wzaWtkkyJ14dTQpqfd)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k10_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Let $k7_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k10_xboolean X0 (k7_xboolean X0 X1) = k3_xboolean X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k3_xboolean (k10_xboolean X0 X1) = k10_xboolean (k3_xboolean X0) X1)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (k3_xboolean (k3_xboolean X0) = X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k7_xboolean X0 X1)) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k10_xboolean (k3_xboolean X0) (k7_xboolean X0 X1) = X1))$$