

t89_xreal_1 (TMdzn-
qzB1imYgixNqbkkgSUTAoqvmHkmA89U)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow ((r1_xxreal_0 X0 k6_numbers) \vee (r1_xxreal_0 (k5_xcmplx_0 X1) (k5_xcmplx_0 X0)))))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k5_xcmplx_0 (k5_xcmplx_0 X0) = X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k5_xcmplx_0 X0)) \wedge (v1_xreal_0 (k5_xcmplx_0 X0))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 (k5_xcmplx_0 X0) (k5_xcmplx_0 X1)) \Rightarrow ((r1_xxreal_0 (k5_xcmplx_0 X0) k6_numbers) \vee (r1_xxreal_0 X1 X0))))$$