

t8_comseq_2
(TMH1f38U11wrSdQrxJs4ZBszk9efFgofJRi)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k2_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_comseq_2 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 X1) \wedge ((\neg v3_xxreal_0 X1) \wedge (\neg v2_xxreal_0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\neg(v1_xboole_0 X0) \wedge ((X0 \neq X1) \wedge (v1_xboole_0 X1)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k17_complex1 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v3_xxreal_0 X1)) \Rightarrow (v3_xxreal_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v2_xxreal_0 X0)) \Rightarrow (v2_xxreal_0 X1))) \quad (5)$$

Assume the following.

$$m1_subset_1 \ k1_xboole_0 \ k4_ordinal1 \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1_funct_1 \ X1) \wedge ((v1_funct_2 \\ & X1 \ k5_numbers \ X0) \wedge (m1_subset_1 \ X1 \ (k1_zfmisc_1 \ (k2_zfmisc_1 \ k5_numbers \\ & X0)))))) \wedge (v7_ordinal1 \ X2)) \Rightarrow (k8_nat_1 \ X0 \ X1 \ X2 = k1_funct_1 \ X1 \ X2) \end{aligned} \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 \ X0) \Rightarrow (k17_complex1 \ X0 = k16_complex1 \ X0) \quad (10)$$

Assume the following.

$$\exists X0. (v1_xboole_0 \ X0) \wedge ((v1_xcmplx_0 \ X0) \wedge ((v1_xxreal_0 \ X0) \wedge (v1_xxreal_0 \ X0))) \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_valued_0 \ X0))) \Rightarrow (v1_xcmplx_0 \ (k1_funct_1 \ X0 \ X1)) \quad (12)$$

Assume the following.

$$v1_xboole_0 \ k1_xboole_0 \quad (13)$$

Assume the following.

$$v1_membered \ k2_numbers \quad (14)$$

Assume the following.

$$m2_subset_1 \ k6_numbers \ k1_numbers \ k5_numbers \quad (15)$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 \ X0) \Rightarrow (v1_xxreal_0 \ (k16_complex1 \ X0)) \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_funct_1 \ X0) \wedge ((v1_funct_2 \ X0 \ k5_numbers \ k2_numbers) \wedge \\ & (m1_subset_1 \ X0 \ (k1_zfmisc_1 \ (k2_zfmisc_1 \ k5_numbers \ k2_numbers)))))) \Rightarrow \\ & ((v1_comseq_2 \ X0) \Leftrightarrow (\exists X1. (m1_subset_1 \ X1 \ k1_numbers) \wedge \\ & \forall X2. (m2_subset_1 \ X2 \ k1_numbers \ k5_numbers) \Rightarrow (\neg r1_xxreal_0 \\ & X1 \ (k17_complex1 \ (k8_nat_1 \ k2_numbers \ X0 \ X2)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (18)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0)\wedge(v3_xxreal_0 X0))\Rightarrow((\neg v1_xboole_0 X0)\wedge((v1_xxreal_0 X0)\wedge(\neg v2_xxreal_0 X0))) \quad (19)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0)\wedge(v2_xxreal_0 X0))\Rightarrow((\neg v1_xboole_0 X0)\wedge((v1_xxreal_0 X0)\wedge(\neg v3_xxreal_0 X0))) \quad (20)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (21)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.(v1_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_valued_0 X2)) \quad (23)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_funct_1 X0)\wedge((v1_funct_2 X0 k5_numbers k2_numbers)\wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers))))))\Rightarrow \\ & ((v1_comseq_2 X0)\Leftrightarrow(\exists X1.(m1_subset_1 X1 k1_numbers)\wedge(\\ & (\neg r1_xxreal_0 X1 k6_numbers)\wedge(\forall X2.(m2_subset_1 X2 k1_numbers \\ & k5_numbers)\Rightarrow(\neg r1_xxreal_0 X1 (k17_complex1 (k8_nat_1 k2_numbers \\ & X0 X2))))))) \end{aligned}$$