

t8_euclid (TMadKrKucZhHGZDgRabaCK- qZQJJ9ZocM5Xx)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k5_euclid : \iota \Rightarrow \iota$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $k13_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k12_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_square_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_euclid : \iota \Rightarrow \iota$ be given. Let $k39_valued_1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v2_valued_0 : \iota \Rightarrow o$ be given. Let $v4_valued_0 : \iota \Rightarrow o$ be given. Let $v5_valued_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m2_finseq_2 X1 k1_numbers \\ (k4_finseq_2 X0 k1_numbers)) \Rightarrow ((k18_rvsum_1 (k13_rvsum_1 X0 X1) = \\ k6_numbers) \Rightarrow (X1 = k5_finseq_2 k5_numbers X0 k6_numbers))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 \\ X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (r1_xxreal_0 k6_numbers (k18_rvsum_1 \\ (k12_rvsum_1 X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \wedge (k6_square_1 X0 = k6_numbers)) \Rightarrow (X0 = k6_numbers)) \quad (4)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (5)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0) \Rightarrow (\forall X2.(m2_finseq_2 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k7_square_1 X0 = k6_square_1 X0) \quad (9)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge ((v7_ordinal1 X1) \wedge (m1_subset_1 X2 X0))) \Rightarrow (k5_finseq_2 X0 X1 X2 = k2_finseq_2 X1 X2) \quad (11)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k5_euclid X0 = k4_euclid X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (m1_subset_1 X1 (k4_finseq_2 X0 k1_numbers))) \Rightarrow (k13_rvsum_1 X0 X1 = k39_valued_1 X1) \quad (13)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (k12_rvsum_1 X0 = k39_valued_1 X0) \quad (14)$$

Assume the following.

$$\begin{aligned} \exists X0.(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ k5_numbers))) \wedge ((\neg v1_xboole_0 X0) \wedge ((v1_relat_1 X0) \wedge ((v4_relat_1 \\ X0 k5_numbers) \wedge ((v5_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge \\ ((v1_partfun1 X0 k5_numbers) \wedge ((v1_funct_2 X0 k5_numbers k5_numbers) \wedge \\ ((v1_valued_0 X0) \wedge ((v2_valued_0 X0) \wedge ((v3_valued_0 X0) \wedge ((v4_valued_0 \\ X0) \wedge (v5_valued_0 X0))))))))))))) \end{aligned} \quad (15)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0) \Rightarrow (\forall X2.(m2_finseq_2 \\ X2 X0 X1) \Rightarrow (m2_finseq_1 X2 X0)) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ X0)))))) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (\\ (v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \quad (19)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (m1_finseq_2 (k1_euclid X0) k1_numbers) \quad (20)$$

Assume the following.

$$\forall X0.(m1_finseq_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k18_rvsum_1 \\ X0) k1_numbers) \quad (21)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 \\ X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (m2_finseq_1 (k12_rvsum_1 X0) k1_numbers) \quad (22)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 \\ X0) \wedge (v3_valued_0 X0)))) \Rightarrow (k12_euclid X0 = k7_square_1 (k18_rvsum_1 \\ (k12_rvsum_1 X0))) \quad (23)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(k4_euclid\ X0 = k5_finseq_2\ k1_numbers\ X0\ k6_numbers) \quad (24)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(k1_euclid\ X0 = k4_finseq_2\ X0\ k1_numbers) \quad (25)$$

Assume the following.

$$\forall X0.\forall X1.(v1_xboole_0\ X0)\Rightarrow(\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow(v1_xboole_0\ X2)) \quad (26)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1\ X1\ X0)\Rightarrow(v5_relat_1\ X1\ X0) \quad (27)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge(v5_relat_1\ X0\ k1_numbers))\Rightarrow((v1_relat_1\ X0)\wedge(v3_valued_0\ X0)) \quad (28)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers)\Rightarrow(v1_xreal_0\ X0) \quad (29)$$

Theorem 1

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(\forall X1.(m2_finseq_2\ X1\ k1_numbers\ (k1_euclid\ X0))\Rightarrow((k12_euclid\ X1 = k6_numbers)\Rightarrow(X1 = k5_euclid\ X0)))$$