

t8_euclid_2

(TMaorA6gMofDkZ6Y1c7vEAUPpskPkrepSPk)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k23_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_square_1 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_square_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 X0) \wedge (v3_valued_0 X0)))) \Rightarrow (r1_xxreal_0 k6_numbers (k12_euclid X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (\neg (k3_xcmplx_0 X0 X1 = k6_numbers) \wedge ((X0 \neq k6_numbers) \wedge (X1 \neq k6_numbers)))) \quad (2)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 X0) \wedge (v3_valued_0 X0)))) \Rightarrow (k5_square_1 (k12_euclid X0) = k23_rvsum_1 X0 X0) \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k5_square_1 X0 = k3_square_1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xreal_0 (k6_square_1 X0)) \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 X0) \wedge (v3_valued_0 X0)))) \Rightarrow (m1_subset_1 (k12_euclid X0) k1_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \Rightarrow (\forall X1. (v1_xreal_0 X1) \Rightarrow ((X1 = k6_square_1 X0) \Leftrightarrow ((r1_xxreal_0 k6_numbers X1) \wedge (k3_square_1 X1 = X0))))) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_square_1 X0 = k3_xcmplx_0 X0 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 X0) \wedge (v3_valued_0 X0)))) \Rightarrow ((k23_rvsum_1 X0 X0 = k6_numbers) \Leftrightarrow (k12_euclid X0 = k6_numbers))$$