

# t8\_euclid\_5

## (TMHK8X5RXF6dKRKY1z8z3kZ85vYrAskFzXd)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k1\_rlvect\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_3 : \iota$  be given. Let  $k4\_euclid\_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_euclid\_5 : \iota \Rightarrow \iota$  be given. Let  $k2\_euclid\_5 : \iota \Rightarrow \iota$  be given. Let  $k3\_euclid\_5 : \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 \\ & X1 (u1\_struct\_0 (k15\_euclid np\_3))) \Rightarrow (k1\_rlvect\_1 (k15\_euclid \\ & np\_3) X1 X0 = k4\_euclid\_5 (k8\_real\_1 X0 (k1\_euclid\_5 X1)) (k8\_real\_1 \\ & X0 (k2\_euclid\_5 X1)) (k8\_real\_1 X0 (k3\_euclid\_5 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 \\ & X1 k1\_numbers) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_numbers) \Rightarrow ((k1\_euclid\_5 \\ & (k4\_euclid\_5 X0 X1 X2) = X0) \wedge ((k2\_euclid\_5 (k4\_euclid\_5 X0 X1 X2) = \\ & X1) \wedge (k3\_euclid\_5 (k4\_euclid\_5 X0 X1 X2) = X2)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1\_xreal\_0 X0) \wedge ((v1\_xreal\_0 \\ & X1) \wedge (v1\_xreal\_0 X2))) \Rightarrow (m1\_subset\_1 (k4\_euclid\_5 X0 X1 X2) (u1\_struct\_0 \\ & (k15\_euclid np\_3))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (4)$$

### Theorem 1

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 \\ & X1 k1\_numbers) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_numbers) \Rightarrow (\forall X3. \\ & (m1\_subset\_1 X3 k1\_numbers) \Rightarrow (k1\_rlvect\_1 (k15\_euclid np\_3) \\ & (k4\_euclid\_5 X1 X2 X3) X0 = k4\_euclid\_5 (k8\_real\_1 X0 X1) (k8\_real\_1 \\ & X0 X2) (k8\_real\_1 X0 X3)))) \end{aligned}$$