

t8_finseq_2 (TM-
SuXRQxRx5aRBaNn33FWd7762pSse7wbZ9)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X0 \in k2_finseq_1 X1) \Leftrightarrow ((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 X0 X1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2.(v7_ordinal1 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 X0 (k2_xcmplx_0 X1 X2))))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (3)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2.(v7_ordinal1 X2) \Rightarrow ((X0 \in k2_finseq_1 X1) \Rightarrow (X0 \in k2_finseq_1 (k2_xcmplx_0 X1 X2)))))$$