

t8_flang_3

(TMJTy4QCdWTHspirSxqWt7uETPVQegCnuy9)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_flang_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 \\ & \quad X3) \Rightarrow ((r1_xxreal_0 X2 (k2_xcmplx_0 X3 np_1)) \Rightarrow (k4_subset_1 (k8_afinsq_1 \\ & \quad X0) (k1_flang_2 X0 X1 X2 X3) (k1_flang_3 X0 X1 (k2_xcmplx_0 X3 np_1)) = \\ & \quad k1_flang_3 X0 X1 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (k1_flang_2 X0 X1 X2 X2 = k7_flang_1 \\ & \quad X0 X1 X2)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow (r1_xxreal_0 \\ & \quad X0 (k2_xcmplx_0 X0 X1))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \tag{4}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{5}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & X0)))\Rightarrow(\forall X2.(v7_ordinal1 X2)\Rightarrow(k4_subset_1 (k8_afinsq_1 \\ X0) (k7_flang_1 X0 X1 X2) (k1_flang_3 X0 X1 (k2_xcmplx_0 X2 np_1)) = \\ & k1_flang_3 X0 X1 X2)) \end{aligned}$$