

t8_interva1
(TMVDdfPeS2FNTkpZU8YJbLk6rRt7U92NcFu)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_interval : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (\forall X3. (X3 \in k1_interval \\ & X0 X1 X2) \Leftrightarrow ((r1_tarski X1 X3) \wedge (r1_tarski X3 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (X0 = X1) \Leftrightarrow ((r1_tarski X0 X1) \wedge (r1_tarski X1 X0)) \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k1_interval X0 X1 X1 = k1_tarski X1)$$