

t8_lexbfs

(TML3A6jFUfsWCMqaDogWCEag2wrcm4U5Fxb)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_lexbfs : \iota \Rightarrow o$ be given. Let $v13_glib_000 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \quad (1)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (m1_subset_1 X1 k5_numbers)) \Rightarrow (k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge (v1_partfun1 X0 k5_numbers)))) \Rightarrow ((v3_lexbfs X0) \Leftrightarrow (\exists X1.(v7_ordinal1 X1) \wedge (\forall X2.(v7_ordinal1 X2) \Rightarrow ((r1_xxreal_0 X1 X2) \Rightarrow (k1_funct_1 X0 X1 = k1_funct_1 X0 X2))))) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge (v1_partfun1 X0 k5_numbers)))) \Rightarrow ((v13_glib_000 X0) \Leftrightarrow (\exists X1. \\ (v7_ordinal1 X1) \wedge (k1_funct_1 X0 X1 = k1_funct_1 X0 (k1_nat_1 X1 \\ np_1)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (9)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (v1_xxreal_0 X0) \quad (10)$$

Theorem 1

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge (v1_partfun1 X0 k5_numbers)))) \Rightarrow ((v3_lexbfs X0) \Rightarrow (v13_glib_000 \\ X0)) \end{aligned}$$