

t8\_midsp\_3 (TMXC-  
sWC2aCoEZQVv5w8RzqDqoqx4zqe4cwH)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_midsp\_3 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\forall X3.(v1\_xreal\_0 X3) \Rightarrow (((r1\_xxreal\_0 \\ & X0 X1) \wedge (r1\_xxreal\_0 X2 X3)) \Rightarrow (r1\_xxreal\_0 (k2\_xcmplx\_0 X0 X2) ( \\ & k2\_xcmplx\_0 X1 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\neg(X0 \in X1) \wedge ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X2)) \wedge (v1\_xboole\_0 X2)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (r1\_xxreal\_0 X0 (k2\_xcmplx\_0 X0 X1))) \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$r1\_xxreal\_0 np\_1 np\_1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge((\neg v1\_xboole\_0 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))))\Rightarrow(\forall X2.(m2\_subset\_1 X2 X0 X1)\Leftrightarrow(m1\_subset\_1 X2 X1)) \quad (7)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (9)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (10)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(m2\_subset\_1 (k2\_nat\_1 X0 X1) k1\_numbers k5\_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k5\_numbers)\Rightarrow(\forall X1.(m1\_subset\_1 X1 k5\_numbers)\Rightarrow((m1\_midsp\_3 X1 X0)\Leftrightarrow((r1\_xxreal\_0 np\_1 X1)\wedge(r1\_xxreal\_0 X1 (k2\_nat\_1 X0 np\_1)))))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_nat\_1 X1 X0) \quad (14)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (15)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(v1\_xreal\_0 X0) \quad (16)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 k5\_numbers)\Rightarrow(\forall X1.(m1\_subset\_1 X1 k5\_numbers)\Rightarrow((r1\_xxreal\_0 X0 X1)\Rightarrow(m1\_midsp\_3 (k2\_nat\_1 X0 np\_1) X1)))$$