

t8_modelc_2

(TMUxziUNds2suUCKLrPBhhpyfLE8oPrzXRG)

October 27, 2020

Let $v1_modelc_2 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v7_modelc_2 : \iota \Rightarrow o$ be given. Let $k7_modelc_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_modelc_2 : \iota \Rightarrow \iota$ be given. Let $k12_modelc_2 : \iota \Rightarrow \iota$ be given. Let $v4_modelc_2 : \iota \Rightarrow o$ be given. Let $v5_modelc_2 : \iota \Rightarrow o$ be given. Let $v8_modelc_2 : \iota \Rightarrow o$ be given. Let $k4_modelc_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_modelc_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_modelc_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.((v1_modelc_2 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\\
 & (\neg(\neg v4_modelc_2 X0) \wedge (\neg v5_modelc_2 X0) \wedge (\neg v7_modelc_2 X0) \wedge \\
 & (\neg v8_modelc_2 X0))) \Rightarrow (\forall X1.((v1_modelc_2 X1) \wedge (m2_finseq_1 \\
 & X1 k5_numbers)) \Rightarrow (((v4_modelc_2 X0) \Rightarrow ((X1 = k12_modelc_2 X0) \Leftrightarrow (\\
 & \exists X2.((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge (\\
 & k4_modelc_2 X2 X1 = X0)))) \wedge (((v5_modelc_2 X0) \Rightarrow ((X1 = k12_modelc_2 \\
 & X0) \Leftrightarrow (\exists X2.((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge \\
 & (k5_modelc_2 X2 X1 = X0)))) \wedge (((v7_modelc_2 X0) \Rightarrow ((X1 = k12_modelc_2 \\
 & X0) \Leftrightarrow (\exists X2.((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge \\
 & (k7_modelc_2 X2 X1 = X0)))) \wedge (\neg(\neg v4_modelc_2 X0) \wedge (\neg v5_modelc_2 \\
 & X0) \wedge (\neg v7_modelc_2 X0) \wedge (\neg(X1 = k12_modelc_2 X0) \Leftrightarrow (\exists X2. \\
 & ((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge (k8_modelc_2 \\
 & X2 X1 = X0))))))))))
 \end{aligned}
 \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_modelc_2 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\\
& (\neg(\neg v4_modelc_2 X0) \wedge (\neg v5_modelc_2 X0) \wedge (\neg v7_modelc_2 X0) \wedge \\
& (\neg v8_modelc_2 X0))) \Rightarrow (\forall X1.((v1_modelc_2 X1) \wedge (m2_finseq_1 \\
& X1 k5_numbers)) \Rightarrow (((v4_modelc_2 X0) \Rightarrow ((X1 = k11_modelc_2 X0) \Leftrightarrow (\\
& \exists X2.((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge (\\
& k4_modelc_2 X1 X2 = X0)))) \wedge ((v5_modelc_2 X0) \Rightarrow ((X1 = k11_modelc_2 \\
& X0) \Leftrightarrow (\exists X2.((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge \\
& (k5_modelc_2 X1 X2 = X0)))) \wedge ((v7_modelc_2 X0) \Rightarrow ((X1 = k11_modelc_2 \\
& X0) \Leftrightarrow (\exists X2.((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge \\
& (k7_modelc_2 X1 X2 = X0)))) \wedge (\neg(\neg v4_modelc_2 X0) \wedge (\neg v5_modelc_2 \\
& X0) \wedge (\neg v7_modelc_2 X0) \wedge (\neg(X1 = k11_modelc_2 X0) \Leftrightarrow (\exists X2. \\
& ((v1_modelc_2 X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge (k8_modelc_2 \\
& X1 X2 = X0)))))))))
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_modelc_2 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\\
& (v7_modelc_2 X0) \Leftrightarrow (\exists X1.((v1_modelc_2 X1) \wedge (m2_finseq_1 \\
& X1 k5_numbers)) \wedge (\exists X2.((v1_modelc_2 X2) \wedge (m2_finseq_1 \\
& X2 k5_numbers)) \wedge (X0 = k7_modelc_2 X1 X2)))
\end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v1_modelc_2 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\\
& (v7_modelc_2 X0) \Rightarrow (X0 = k7_modelc_2 (k11_modelc_2 X0) (k12_modelc_2 \\
& X0)))
\end{aligned}$$