

t8_polyform

(TMVm9ivusdfeUgio79JRFi3yJzRgGdnBW2g)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_abian : \iota \Rightarrow o$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k3_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k3_power X0 X1 = k1_newton X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0.((v1_int_1 X0) \wedge (v1_abian X0)) \Rightarrow (k3_power (k4_xcmplx_0 np_1) X0 = np_1) \quad (2)$$

Assume the following.

$$((v2_xreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (v1_xreal_0 (k4_xcmplx_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_int_1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((v1_abian X0) \Rightarrow (k1_newton (k4_xcmplx_0 np_1) X0 = np_1))$$