

t91_glib_000

(TMT6spL1Vy8T6wAh3nLxNMgRTeT2DSpfMfb)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $r5_glib_000 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_glib_000 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.((v1_relat_1 \\ X1) \wedge ((v4_relat_1 X1 k5_numbers) \wedge ((v1_funct_1 X1) \wedge ((v1_finset_1 \\ X1) \wedge (v1_glib_000 X1)))))) \Rightarrow ((r5_glib_000 X0 X1) \Leftrightarrow ((m1_glib_000 \\ X0 X1) \wedge (m1_glib_000 X1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.(m1_glib_000 \\ X1 X0) \Rightarrow (\forall X2.(m1_glib_000 X2 X1) \Rightarrow (m1_glib_000 X2 X0))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.((v1_relat_1 \\ X1) \wedge ((v4_relat_1 X1 k5_numbers) \wedge ((v1_funct_1 X1) \wedge ((v1_finset_1 \\ X1) \wedge (v1_glib_000 X1)))))) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge ((v4_relat_1 \\ X2 k5_numbers) \wedge ((v1_funct_1 X2) \wedge ((v1_finset_1 X2) \wedge (v1_glib_000 \\ X2)))))) \Rightarrow (((r5_glib_000 X0 X1) \wedge (m1_glib_000 X2 X0)) \Rightarrow (m1_glib_000 \\ X2 X1))) \end{aligned}$$