

t92_mesfunc6 (TMMcH- fKhPb6TPbRTtS2RXKDe7n2Zx1WzCeQ)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r3_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_supinf_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
& \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))))) \Rightarrow \\
& (\forall X3. ((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k7_numbers)))) \Rightarrow (\forall X4. (m2_subset_1 X4 (k1_zfmisc_1 X0) \\
& X1) \Rightarrow (\forall X5. (m2_subset_1 X5 (k1_zfmisc_1 X0) X1) \Rightarrow (((r1_mesfunc5 \\
& X0 X1 X2 X3) \wedge (r1_xboole_0 X4 X5)) \Rightarrow (k7_mesfunc5 X0 X1 X2 (k2_partfun1 \\
& X0 k7_numbers X3 (k6_prob_1 X0 X1 X4 X5)) = k3_supinf_2 (k7_mesfunc5 \\
& X0 X1 X2 (k2_partfun1 X0 k7_numbers X3 X4)) (k7_mesfunc5 X0 X1 X2 (\\
& k2_partfun1 X0 k7_numbers X3 X5))))))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X2) \wedge \\
& (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow (k2_partfun1 \\
& X0 X1 X2 X3 = k5_relat_1 X2 X3)
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 \\ & (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) \\ & (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow \\ & ((v1_funct_1 (k1_mesfunc5 X0 X1))\wedge(m1_subset_1 (k1_mesfunc5 \\ & X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow(\\ & k1_mesfunc5 X0 X1 = X1) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge \\ & ((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 \\ & X2 X1 k7_numbers)\wedge((v10_valued_0 X2)\wedge((v6_supinf_2 X2)\wedge((v4_measure1 \\ & X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers))))))\Rightarrow \\ & (\forall X3.((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k1_numbers))))\Rightarrow((r3_mesfunc6 X0 X1 X2 X3)\Leftrightarrow(r1_mesfunc5 X0 X1 \\ & X2 (k1_mesfunc5 X0 X3)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge \\ & ((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 \\ & X2 X1 k7_numbers)\wedge((v10_valued_0 X2)\wedge((v6_supinf_2 X2)\wedge((v4_measure1 \\ & X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers))))))\Rightarrow \\ & (\forall X3.((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k1_numbers))))\Rightarrow(k1_mesfunc6 X0 X1 X2 X3 = k7_mesfunc5 X0 X1 X2 \\ & (k1_mesfunc5 X0 X3)))))) \end{aligned} \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\ & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\ & (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k1_numbers)))) \Rightarrow (\forall X4.(m2_subset_1 X4 (k1_zfmisc_1 X0) \\ & X1) \Rightarrow (\forall X5.(m2_subset_1 X5 (k1_zfmisc_1 X0) X1) \Rightarrow (((r3_mesfunc6 \\ & X0 X1 X2 X3) \wedge (r1_xboole_0 X4 X5)) \Rightarrow (k1_mesfunc6 X0 X1 X2 (k2_partfun1 \\ & X0 k1_numbers X3 (k6_prob_1 X0 X1 X4 X5)) = k3_supinf_2 (k1_mesfunc6 \\ & X0 X1 X2 (k2_partfun1 X0 k1_numbers X3 X4)) (k1_mesfunc6 X0 X1 X2 (\\ & k2_partfun1 X0 k1_numbers X3 X5)))))))))) \end{aligned}$$