

t92_xcplx_1
(TMb47yJX88wTuNDbpU9qcxGVJus7Kx7rcR7)

October 27, 2020

Let $v1_xcplx_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (\forall X2. \\ & (v1_xcplx_0 X2) \Rightarrow (k7_xcplx_0 X0 (k3_xcplx_0 X1 X2) = k7_xcplx_0 \\ & (k7_xcplx_0 X0 X1) X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow ((\\ & X0 \neq k6_numbers) \Rightarrow (k3_xcplx_0 (k7_xcplx_0 X1 X0) X0 = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xcplx_0 X0) \wedge (v1_xcplx_0 X1)) \Rightarrow (\\ & v1_xcplx_0 (k7_xcplx_0 X0 X1)) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (\forall X2. \\ & (v1_xcplx_0 X2) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (k7_xcplx_0 X1 X2 = k3_xcplx_0 \\ & (k7_xcplx_0 X1 (k3_xcplx_0 X2 X0) X0)))) \end{aligned}$$