

t93_xxreal_1

(TMGqUEzfl1ykvQNvvnwWWHpvLnNG3hW3T9Dv1)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_xxreal_0 \\ & X0 X1) \Rightarrow (r1_xboole_0 (k1_xxreal_1 X2 X0) (k4_xxreal_1 X1 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2.(v1_xxreal_0 \\ & X2) \Rightarrow (\neg(X0 \in k1_xxreal_1 X1 X2) \wedge ((\neg X0 \in k4_xxreal_1 X1 X2) \wedge ((X0 \neq \\ & X1) \wedge (X0 \neq X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_tarski (\\ & k4_xxreal_1 X0 X1) (k1_xxreal_1 X2 X3)) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (\\ & (r1_xxreal_0 X2 X0) \wedge (r1_xxreal_0 X1 X3)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(\neg(\neg r1_xboole_0 X0 X1) \wedge (\forall X2.\neg(X2 \in \\ & X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2.(X2 \in X0) \wedge (X2 \in X1)) \wedge (r1_xboole_0 \\ & X0 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (r1_tarski \\ & (k4_xxreal_1 X0 X1) (k1_xxreal_1 X0 X1))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k1_xxreal_1 X1 X2) \Leftrightarrow ((r1_xxreal_0 X1 X0) \wedge \\ & (r1_xxreal_0 X0 X2)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. (v1_xxreal_0 \\ & X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Rightarrow ((X0 \in k1_xxreal_1 X1 X2) \wedge ((X0 \neq X1) \wedge \\ & (X0 \neq X2)))))) \end{aligned} \tag{7}$$

Theorem 1

$$\begin{aligned} & \forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3. (v1_xxreal_0 X3) \Rightarrow ((r1_xxreal_0 \\ & X0 X1) \Rightarrow (r1_xboole_0 (k4_xxreal_1 X2 X0) (k1_xxreal_1 X1 X3)))))) \end{aligned}$$