

t93\_xxreal\_2

(TMKZDT5bTbLc2wZBwusPYuhC2ZYuVwE3UiK)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xxreal\_0 : \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow ((r1\_xxreal\_0 k1\_xxreal\_0 X0) \Rightarrow (X0 = k1\_xxreal\_0)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.\neg(X0 \neq k1\_tarski X1) \wedge (\forall X2.(m1\_subset\_1 X2 X0) \Rightarrow (X2 = X1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k7\_numbers) \Rightarrow (v1\_xxreal\_0 X0) \quad (5)$$

**Theorem 1**

$$\forall X0.((\neg v1\_xboole\_0 X0) \wedge (m1\_subset\_1 X0 (k1\_zfmisc\_1 k7\_numbers))) \Rightarrow ((\forall X1.(m1\_subset\_1 X1 k7\_numbers) \Rightarrow ((X1 \in X0) \Rightarrow (r1\_xxreal\_0 k1\_xxreal\_0 X1))) \Rightarrow (X0 = k1\_tarski k1\_xxreal\_0))$$