

t93_zf_lang1
(TMPPT1LzPBAX8VLLL1Mqwnp8R4outiiB3L7)

October 27, 2020

Let $v1_zf_lang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zf_lang : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $r2_zf_model : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_zf_model : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zf_lang1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & (m2_subset_1 X1 k5_numbers k1_zf_lang) \Rightarrow (\forall X2.(\neg v1_xboole_0 \\ & X2) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge ((v1_funct_2 X3 k1_zf_lang \\ & X2) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 k1_zf_lang X2)))))) \Rightarrow \\ & ((r1_zf_model X2 X3 (k13_zf_lang X1 X0)) \Leftrightarrow (\exists X4.(m1_subset_1 \\ & X4 X2) \wedge (r1_zf_model X2 (k2_zf_lang1 X2 X3 X1 X4) X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\neg v1_xboole_0 k1_zf_lang \quad (4)$$

Assume the following.

$$m1_subset_1 k1_zf_lang (k1_zfmisc_1 k5_numbers) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((m1_subset_1 X0 k1_zf_lang) \wedge ((v1_zf_lang \\ & X1) \wedge (m1_finseq_1 X1 k5_numbers))) \Rightarrow ((v1_zf_lang (k13_zf_lang \\ & X0 X1)) \wedge (m2_finseq_1 (k13_zf_lang X0 X1) k5_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_zf_lang X1) \wedge \\ m2_finseq_1 X1 k5_numbers)) \Rightarrow ((r2_zf_model X0 X1) \Leftrightarrow (\forall X2. \\ ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 k1_zf_lang X0) \wedge (m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 k1_zf_lang X0)))))) \Rightarrow (r1_zf_model \\ X0 X2 X1))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (8)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ (m2_subset_1 X1 k5_numbers k1_zf_lang) \Rightarrow (\forall X2.(\neg v1_xboole_0 \\ X2) \Rightarrow ((r2_zf_model X2 (k13_zf_lang X1 X0)) \Leftrightarrow (\forall X3.((v1_funct_1 \\ X3) \wedge ((v1_funct_2 X3 k1_zf_lang X2) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\ (k2_zfmisc_1 k1_zf_lang X2)))))) \Rightarrow (\exists X4.(m1_subset_1 X4 \\ X2) \wedge (r1_zf_model X2 (k2_zf_lang1 X2 X3 X1 X4) X0)))))) \end{aligned}$$