

t94_mesfunc6

(TMEnxPxDvaMp6w2JPfg9sxsSbSMypHPJS3jj)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r3_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k56_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & \quad m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ & \quad r2_relset_1 X0 k7_numbers (k10_mesfunc1 X0 (k1_mesfunc5 X0 X1)) \\ & \quad (k1_mesfunc5 X0 (k56_valued_1 X0 k1_numbers X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\ & \quad ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & \quad \quad (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & \quad X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\ & \quad X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\ & \quad (\forall X3. ((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & \quad \quad X0 k7_numbers)))) \Rightarrow ((\exists X4. (m2_subset_1 X4 (k1_zfmisc_1 \\ & \quad X0) X1) \wedge ((X4 = k1_relset_1 X0 X3) \wedge (r1_mesfunc1 X0 X1 X3 X4))) \Rightarrow ((\\ & \quad r1_mesfunc5 X0 X1 X2 X3) \Leftrightarrow (r1_mesfunc5 X0 X1 X2 (k10_mesfunc1 X0 X3)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((r2_relset_1 X0 X1 X2 X3)\Leftrightarrow(X2 = X3)) \quad (3)$$

Assume the following.

$$v3_membered k1_numbers \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_membered X1)\wedge((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 (k56_valued_1 X0 X1 X2))\wedge(m1_subset_1 (k56_valued_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow((v1_funct_1 (k1_mesfunc5 X0 X1))\wedge(m1_subset_1 (k1_mesfunc5 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0)\wedge((v5_relat_1 X1 k7_numbers)\wedge(v1_funct_1 X1))))\Rightarrow((v1_funct_1 (k10_mesfunc1 X0 X1))\wedge(m1_subset_1 (k10_mesfunc1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \quad (7)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow(k1_mesfunc5 X0 X1 = X1)) \quad (8)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 X2 X1 k7_numbers)\wedge((v10_valued_0 X2)\wedge((v6_supinf_2 X2)\wedge((v4_measure1 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers))))))))\Rightarrow(\forall X3.((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow((r3_mesfunc6 X0 X1 X2 X3)\Leftrightarrow(r1_mesfunc5 X0 X1 X2 (k1_mesfunc5 X0 X3)))))) \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\forall X3.(\\ & m2_subset_1 X3 (k1_zfmisc_1 X0) X1) \Rightarrow ((r1_mesfunc6 X0 X1 X2 X3) \Leftrightarrow \\ & (r1_mesfunc1 X0 X1 (k1_mesfunc5 X0 X2) X3)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.(v3_membered X0) \Rightarrow (v1_membered X0) \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \quad (13)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\ & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\ & (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k1_numbers)))) \Rightarrow ((\exists X4.(m2_subset_1 X4 (k1_zfmisc_1 \\ & X0) X1) \wedge ((X4 = k1_rset_1 X0 X3) \wedge (r1_mesfunc6 X0 X1 X3 X4))) \Rightarrow ((\\ & r3_mesfunc6 X0 X1 X2 X3) \Leftrightarrow (r3_mesfunc6 X0 X1 X2 (k56_valued_1 X0 k1_numbers \\ & X3))))))))) \end{aligned}$$