

t94_relat_1 (TM-
FZTCh9yevkQSimbm5oWAfbe7LRCVv69VX)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k6_relat_1 X0 X1 = k3_relat_1 X1 (k4_relat_1 X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski (k10_xtuple_0 X1) X0) \Rightarrow (k3_relat_1 X1 (k4_relat_1 X0) = X1)) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski (k10_xtuple_0 X1) X0) \Rightarrow (k6_relat_1 X0 X1 = X1))$$