

t94_xcplx_1
(TMRjuW7s2mgeR54ipX1cdNoamFsw6Xcg9y7)

October 27, 2020

Let $v1_xcplx_0 : \iota \Rightarrow o$ be given. Let $k3_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (X1 = k7_xcplx_0 (k3_xcplx_0 X1 X0) X0))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (\forall X2.(v1_xcplx_0 X2) \Rightarrow (k7_xcplx_0 (k7_xcplx_0 X0 X1) X2 = k7_xcplx_0 (k7_xcplx_0 X0 X2) X1))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcplx_0 X0) \wedge (v1_xcplx_0 X1)) \Rightarrow (v1_xcplx_0 (k3_xcplx_0 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcplx_0 X0) \wedge (v1_xcplx_0 X1)) \Rightarrow (k3_xcplx_0 X0 X1 = k3_xcplx_0 X1 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (\forall X2.(v1_xcplx_0 X2) \Rightarrow (\forall X3.(v1_xcplx_0 X3) \Rightarrow ((k3_xcplx_0 X2 X0 = k3_xcplx_0 X3 X1) \Rightarrow ((X0 = k6_numbers) \vee ((X1 = k6_numbers) \vee (k7_xcplx_0 X2 X1 = k7_xcplx_0 X3 X0)))))))) \quad (5)$$