

t96_clvect_2

(TMTKY7vn7Bc8RnayzSiMR7FLHekiXkT88Jk)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v2_clvect_1 : \iota \Rightarrow o$ be given. Let $v3_clvect_1 : \iota \Rightarrow o$ be given. Let $v4_clvect_1 : \iota \Rightarrow o$ be given. Let $v5_clvect_1 : \iota \Rightarrow o$ be given. Let $v2_csspace : \iota \Rightarrow o$ be given. Let $l1_csspace : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v3_funct_1 : \iota \Rightarrow o$ be given. Let $k1_valued_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_valued_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_valued_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k9_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Let $l1_clvect_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers X0)))))) \Rightarrow (\forall X2. (m2_valued_0 X2 X0 X1) \Leftrightarrow (m1_valued_0 \\ & X2 X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{2}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0))))))\wedge(v7_ordinal1 X2)))\Rightarrow(k1_valued_0 X0 X1 X2 = k9_nat_1 X1 X2) \quad (4)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (5)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0)\wedge(l1_struct_0 X0))\Rightarrow(\neg v1_xboole_0 (u1_struct_0 X0)) \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (7)$$

Assume the following.

$$\forall X0.(l2_algstr_0 X0)\Rightarrow((l2_struct_0 X0)\wedge(l1_algstr_0 X0)) \quad (8)$$

Assume the following.

$$\forall X0.(l1_csspace X0)\Rightarrow(l1_clvect_1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(l1_clvect_1 X0)\Rightarrow(l2_algstr_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(l1_algstr_0 X0)\Rightarrow(l1_struct_0 X0) \quad (11)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0))))))\wedge(v7_ordinal1 X2)))\Rightarrow(m2_valued_0 (k1_valued_0 X0 X1 X2) X0 X1) \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge((v3_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))))\Rightarrow(\forall X2.(m1_valued_0 X2 X1)\Rightarrow(v3_funct_1 X2)) \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (v13_algstr_0 X0) \wedge (v2_rlvect_1 \\ & X0) \wedge (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v2_clvect_1 X0) \wedge \\ & ((v3_clvect_1 X0) \wedge (v4_clvect_1 X0) \wedge (v5_clvect_1 X0) \wedge (v2_csspace \\ & X0) \wedge (l1_csspace X0)))))) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ & ((v1_funct_2 X1 k5_numbers (u1_struct_0 X0)) \wedge (m1_subset_1 X1 \\ & (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (u1_struct_0 X0)))))) \Rightarrow \\ & (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow ((v3_funct_1 \\ & X1) \Rightarrow (v3_funct_1 (k1_valued_0 (u1_struct_0 X0) X1 X2)))))) \end{aligned}$$