

t96_finseq_6 (TMEonzn-
phe7HEXXDBMAh7ZS5nT3cwKy6M11)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_finseq_6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_r_finseq : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 X0) \Rightarrow (k1_finseq_6 X0 (k2_finseq_4 \\ & X0 X1 X2) X1 = k2_finseq_4 X0 X1 X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 X0) \Rightarrow ((X1 \neq X2) \Rightarrow (k2_finseq_5 X0 (k2_finseq_4 \\ & X0 X1 X2) X2 = k12_finseq_1 X0 X2)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 X0) \Rightarrow ((X1 \neq X2) \Rightarrow (k1_finseq_5 X0 (k2_finseq_4 \\ & X0 X1 X2) X2 = k2_finseq_4 X0 X1 X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 X0) \Rightarrow (k2_r_finseq X0 np_1 (k2_finseq_4 \\ & X0 X1 X2) = k12_finseq_1 X0 X2))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. \forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.k9_finseq_1 X0 = k5_finseq_1 X0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X1 X0)\wedge(m1_finseq_1 X2 X0))\Rightarrow(k8_finseq_1 X0 X1 X2 = k7_finseq_1 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge((m1_subset_1 X1 X0)\wedge(m1_subset_1 X2 X0)))\Rightarrow(k2_finseq_4 X0 X1 X2 = k10_finseq_1 X1 X2) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k10_xtuple_0 (k10_finseq_1 X0 X1) = k2_tarSKI X0 X1 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge((m1_subset_1 X1 X0)\wedge(m1_subset_1 X2 X0)))\Rightarrow(m2_finseq_1 (k2_finseq_4 X0 X1 X2) X0) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.k10_finseq_1 X0 X1 = k7_finseq_1 (k9_finseq_1 X0) (k9_finseq_1 X1) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k2_tarSKI X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 = X0)\vee(X3 = X1))) \quad (14)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m2_finseq_1 X1 X0)\Rightarrow(\forall X2.(m1_subset_1 X2 X0)\Rightarrow(((X2 \in k10_xtuple_0 X1)\Rightarrow(k1_finseq_6 X0 X1 X2 = k8_finseq_1 X0 (k2_finseq_5 X0 X1 X2) (k2_rfinseq X0 np_1 (k1_finseq_5 X0 X1 X2))))\wedge((\neg X2 \in k10_xtuple_0 X1)\Rightarrow(k1_finseq_6 X0 X1 X2 = X1)))))) \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2. (m1_subset_1 X2 X0) \Rightarrow (k1_finseq_6 X0 (k2_finseq_4 \\ & X0 X1 X2) X2 = k2_finseq_4 X0 X2 X2))) \end{aligned}$$