

t96_xboolean (TM- SoK7ArvuDqd6CtXNeSMmsXTPrGMPyWd7F)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k8_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (\forall X2. \\ & (v1_xboolean X2) \Rightarrow (k4_xboolean X0 (k10_xboolean X1 X2) = k10_xboolean \\ & (k4_xboolean X0 X1) (k4_xboolean X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (k3_xboolean (k3_xboolean X0) = X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k10_xboolean X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k7_xboolean X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k4_xboolean X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k8_xboolean X0 X1 = k3_xboolean (k4_xboolean X0 X1))) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k10_xboolean X0 X1 = k3_xboolean (k7_xboolean X0 X1))) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xboolean\ X0) \Rightarrow (\forall X1.(v1_xboolean\ X1) \Rightarrow (\forall X2. \\ & (v1_xboolean\ X2) \Rightarrow (k8_xboolean\ X0\ (k10_xboolean\ X1\ X2) = k7_xboolean \\ & (k4_xboolean\ X0\ X1)\ (k4_xboolean\ X0\ X2)))) \end{aligned}$$