

t97_funct_1
(TMHH4iE1q74GEBR9X2vitLfNHr1wP2my3WW)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (r1_tarski (k9_xtuple_0 (k5_relat_1 X1 X0)) X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k5_relat_1 X2 X1)) \Rightarrow (k1_funct_1 (k5_relat_1 X2 X1) X0 = k1_funct_1 X2 X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_relat_1 (k5_relat_1 X0 X1)) \wedge (v1_funct_1 (k5_relat_1 X0 X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X0) \Rightarrow (v1_relat_1 (k5_relat_1 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (5)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. (X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (7)$$

Theorem 1

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v1_funct_1 X1))\Rightarrow(r1_tarski \\ (k10_xtuple_0 (k5_relat_1 X1 (k1_tarski X0)) (k1_tarski (k1_funct_1 \\ X1 X0)))$$