

t9_afinsq_1
(TMGzu1J8nGYA7s1pxVpW1wpdX3RNbnrhxpz)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v5_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $k1_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v1_card_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X0 \in X1) \Leftrightarrow (\neg r1_xxreal_0 X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((k9_xtuple_0 X0 = k9_xtuple_0 X1) \wedge (\forall X2.(X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 X0 X2 = k1_funct_1 X1 X2)))) \Rightarrow (X0 = X1))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (k2_afinsq_1 X0 = k9_xtuple_0 X0) \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (k1_afinsq_1 X0 = k1_card_1 X0) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (k1_card_1 X0 = k9_xtuple_0 X0) \quad (6)$$

Assume the following.

$$\forall X0.(v1_finset_1 X0) \Rightarrow ((v1_finset_1 (k1_card_1 X0)) \wedge (v1_card_1 (k1_card_1 X0))) \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (m1_subset_1 (k2_afinsq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (k4_tarski X1 X2 \in X0))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (9)$$

Assume the following.

$$\forall X0.((v3_ordinal1 X0) \wedge (v1_finset_1 X0)) \Rightarrow (v7_ordinal1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1_card_1 X0) \Rightarrow (v3_ordinal1 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v6_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v7_ordinal1 X1)) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 k5_numbers)) \Rightarrow (v6_membered X0) \quad (13)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((v5_ordinal1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finset_1 X1)))) \Rightarrow (((k1_afinsq_1 X0 = k1_afinsq_1 X1) \wedge (\forall X2.(v7_ordinal1 X2) \Rightarrow ((\neg r1_xreal_0 (k1_afinsq_1 X0) X2) \Rightarrow (k1_funct_1 X0 X2 = k1_funct_1 X1 X2)))) \Rightarrow (X0 = X1)))$$