

t9_cqc_sim1 (TMHx-
ChdBD3y wzG3sKRHFZsrcGtLTHE6EayW)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m1_subset_1 (k4_finseq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (8)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. (X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (9)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (10)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0)))) \quad (11)$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k10_xtuple_0 X0 = ReplSep (toSet (\lambda X1 : \iota. m1_subset_1 X1 k5_numbers)) (\lambda X1 : \iota. (r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))) (\lambda X1 : \iota. k1_funct_1 X0 X1))$$